

DIGITAL TECHNOLOGY TO CALCULATE WATER FOOTPRINTS FOR DIFFERENT AGRICULTURAL PRODUCTS

Mr. Bhagyesh Shamal Fegade¹, Asst.Prof.Ms.Sapana A.Fegade²

Research Scholar, Department of Computer Applications, Shramsadhana College of Engineering & Technology

Jalgaon.MH, India¹

Assistant Professor, Department of Computer Applications, Shramsadhana College of Engineering & Technology

Jalgaon.MH, India²

Abstract: Agricultural production is inextricably linked with freshwater consumption, rendering quantitative, reliable water-footprint assessments essential for sustainable environmental resource management. This research proposes a robust digital technology-based framework for systematically calculating, validating, organizing, and visualizing the water footprints of diverse agricultural commodities. The proposed digital workflow integrates field data collection, structured product-level digital records, component-wise water-footprint computation engines, automated validation rules, and interactive reporting dashboards. Green (rainwater), blue (surface and groundwater), and grey (pollutant dilution volume) water footprint components are quantified separately and combined to yield precise product-level water footprint indicators (m³/ton). Digital record management drastically improves data traceability by preserving raw data sources, mathematical calculation methods, model version parameters, timestamps, and data quality flags. Operating within a design-oriented engineering framework, this study defines the architectural layers, database schema attributes, sequential data workflows, and a target performance evaluation matrix. The resulting framework provides a transparent foundation for multi-crop comparison, policy reporting, and future integration with Internet of Things (IoT) field sensors, remote sensing, artificial intelligence, and smart agriculture decision-support ecosystems.

Keywords: Water footprint assessment, Digital agriculture, Green-blue-grey water, Data traceability, Precision farming, Smart agriculture, Quality validation, IoT integration.

INTRODUCTION

Water is a fundamental, finite natural resource supporting global agricultural production, food security, and rural livelihoods. Agricultural freshwater consumption varies substantially depending on crop species, regional climate conditions, soil physics, irrigation infrastructure, and agronomic management practices. Water-footprint assessment provides a comprehensive, standardized metric to quantify freshwater consumption across three primary components: green water (soil moisture derived from precipitation consumed via evapotranspiration), blue water (abstracted surface and groundwater utilized for irrigation), and grey water (volume of freshwater required to assimilate pollutant loads to meet water quality standards). Transitioning from traditional manual estimation to a digital technology framework enables systematic, repeatable, and traceable water-footprint accounting.

Need for the Study

Conventional agricultural water-footprint calculations frequently rely on dispersed data sources, inconsistent spatial-temporal resolutions, and manual record keeping, making longitudinal tracking and cross-commodity comparisons highly challenging. A structured digital framework consolidates heterogeneous agricultural datasets into unified, immutable product records. Such a system preserves vital calculation metadata, supports reproducible assessments, and empowers researchers, farmers, water resource managers, and environmental policy analysts with transparent, product-level analytics.

Problem Statement

There is an urgent need for an integrated, transparent digital platform capable of collecting multi-source agricultural data, computing component-wise green, blue, and grey water footprints, enforcing automated data quality validation, and reporting standardized product-level results while preserving full data provenance and auditability.

SCOPE

The scope of this research covers the design, modeling, and architectural specification of a digital framework for collecting, validating, calculating, and visualizing agricultural water footprints. The framework encompasses product-level digital schemas, component-wise calculation engines, data quality flagging, comparative dashboard visualization, and architectural interfaces for future integration with smart-agriculture technologies such as IoT soil moisture sensors, satellite remote sensing, and AI predictive analytics.

OBJECTIVES

The primary objectives of this study are:

- Formulate a standardized digital record structure for agricultural products incorporating metadata, environmental conditions, and production yields.
- Develop an integrated computational engine to separately calculate green, blue, and grey water footprints and combine them into normalized unit metrics (m^3/ton).
- Establish automated validation and quality control protocols to detect missing data, unit discrepancies, and calculation anomalies.
- Design a multi-layered digital platform architecture maintaining end-to-end traceability from raw data inputs to reported outputs.
- Define a comprehensive evaluation framework to benchmark platform performance across completeness, validation accuracy, consistency, traceability, and usability.

REVIEW OF LITERATURE

Mekonnen et al., in (Copernicus 2011) [1], examined about the green, blue, and grey water footprints of crops and derived crop products at a global scale. Their pioneering work provided a foundational accounting methodology demonstrating how freshwater consumption can be rigorously disaggregated across crop types and geographic regions. The study emphasized that disaggregating water use into specific green, blue, and grey components is essential for meaningful product-level comparative evaluation. Their findings establish the necessity for structured, systematic data collection when calculating product-level water footprints in modern digital systems.

Hoekstra et al., in (Earthscan 2011) [2], presented about the Water Footprint Assessment Manual, establishing the global standard principles, definitions, and operational accounting procedures for water footprint assessment. The manual provides a formal structural basis for identifying, quantifying, and interpreting green, blue, and grey water components across agricultural supply chains. Their standardized conceptual framework underpins our proposed digital system design, ensuring that digitized data records strictly align with international water accounting methodologies.

ISO et al., in (ISO 2014) [3], established that ISO 14046 specifies the international principles, requirements, and guidelines for conducting environmental water footprint assessments within life cycle assessment (LCA) frameworks. The standard underscores the mandatory requirement for systematic, transparent, and traceable assessment procedures that reflect environmental contexts. Digital record keeping directly fulfills this requirement by establishing verifiable links between raw field inputs, computational algorithms, and reported footprint metrics.

Mekonnen et al., in (Copernicus 2010) [4], conducted about a high-resolution spatial assessment of the green, blue, and grey water footprint of wheat, proving that agricultural water footprints exhibit extreme spatial and temporal variations based on local climate and agronomic practices. Their spatial modeling highlighted that location-specific soil, weather, and yield parameters are critical for accurate footprint interpretation. This empirical insight directly validates our digital record schema, which explicitly mandates geographic coordinates, seasonal timestamps, and soil metadata.

Allen et al., in (FAO 1998) [5], provided that the FAO Irrigation and Drainage Paper 56 guidelines detail standardized mathematical formulations for estimating crop evapotranspiration (ET_c) and net irrigation requirements using the Penman-Monteith method. These meteorological and agronomic formulas represent core mathematical primitives for calculating green and blue crop water use. Incorporating these standardized FAO formulations into an automated digital calculation layer ensures rigorous mathematical consistency across all product assessments.

Parra-López et al., in (Elsevier 2025) [6], discussed about the rapid evolution and application of digital technologies for agricultural water monitoring, allocation, and decision support. Their comprehensive survey highlighted that digital

tools, cloud databases, and analytical platforms streamline data ingestion and bridge field-level observations with analytical decision systems. Their findings support our proposed digital architecture, demonstrating that digital records transform static water footprint calculations into dynamic, reusable knowledge assets.

Kumar et al., in (Elsevier 2025) [7], explored that the integration of Internet of Things (IoT) wireless sensor networks and artificial intelligence algorithms provides continuous, automated monitoring of agricultural microclimates and soil moisture regimes. Their systematic review demonstrated that intelligent IoT architectures minimize human data entry errors and deliver real-time data feeds. These technological advancements represent a direct future integration pathway for our proposed water footprint digital repository.

Gupta et al., in (Elsevier 2026) [8], highlighted that IoT-enabled smart irrigation systems utilize real-time soil, meteorological, and crop physiological data to optimize precision irrigation scheduling. Their research proved that automated sensing significantly improves water-use efficiency while generating high-frequency digital data logs. Such automated data streams offer ideal input parameters for updating blue and green water footprint calculations dynamically within digital management frameworks.

FAO et al., in (FAO 2011) [9], emphasized that the State of the World's Land and Water Resources for Food and Agriculture demands integrated, multi-source land and water resource management to mitigate systemic environmental risks. The report stressed that sustainable resource governance requires consolidating fragmented agricultural data into coherent analytical frameworks. Our product-level digital record model responds directly to this global directive by harmonizing disparate production data into structured footprint reports.

ISO et al., in (ISO 2017) [10], illustrated that ISO/TR 14073 provides practical illustrative examples and operational guidance for implementing ISO 14046 water footprint assessments in real-world agricultural and industrial scenarios. The technical report demonstrates how mathematical models must be translated into standardized assessment workflows. Our proposed digital architecture reflects these guidelines by embedding explicit data validation, model versioning, and uncertainty flagging into the core software workflow.

Research Gap & Synthesis— Existing literature offers mature mathematical formulations for water footprint calculation (Mekonnen & Hoekstra [1], [4]; Allen et al. [5]) and highlights the rising role of IoT and digital sensing (Parra-López et al. [6]; Gupta et al. [8]). However, there remains a distinct technological gap in establishing a unified, end-to-end product-level digital framework that integrates structured data ingestion, Component-wise (green-blue-grey) calculation, automated quality validation, immutable traceability, and comparative reporting within a single cohesive workflow. The proposed digital platform bridges this operational gap.

METHODOLOGY

This study follows a design-oriented software engineering research methodology. The proposed framework structures the flow of agricultural information through five sequential operational phases: data collection, automated validation, digital record persistence, water-footprint computation, and dashboard reporting, as illustrated in Figure 1 and Figure 3.



Figure 1. Digital Framework for Agricultural Water-Footprint Assessment

Data Sources & Inputs

The framework ingests multi-source data parameters essential for accurate water footprint computation, including commodity identity, geographic coordinates, planting and harvest timestamps, seasonal weather datasets (effective rainfall, reference evapotranspiration), irrigation application volumes, fertilizer/pesticide application rates, and total crop yield.

To enforce standardized data storage, each agricultural product entry is stored as a structured JSON/XML digital object detailing specific metadata and computational attributes:

Digital Schema Field	Attribute Description & Data Requirement
Product & Spatial Identity	Product ID, crop variety, farm location coordinates, regional climate zone, and seasonal period.
Agronomic & Input Parameters	Planting/harvest dates, total crop yield (tons/ha), effective rainfall (mm), irrigation depth (mm), chemical input load (kg/ha).
Calculated Footprint Components	Green water footprint (m ³), Blue water footprint (m ³), Grey water footprint (m ³), Total footprint (m ³).
Normalized Unit Metrics	Water footprint per unit product (m ³ /ton), water productivity ratio (kg/m ³).
Traceability & Quality Metadata	Data source URI, calculation method version, timestamp, validation status flag, and uncertainty score.

Table 1: - Digital Record Structure & Schema Attributes

Water Footprint Calculation Engine

The core computational layer processes green, blue, and grey water components independently before aggregating them into the product-level total, as illustrated in Figure 2. Green water footprint is derived from effective rainfall evapotranspiration over the growing period; blue water footprint represents applied irrigation water volume consumed; grey water footprint is calculated as the volume of freshwater required to dilute chemical leaching to ambient water quality standards. The total water footprint is normalized by crop yield to produce unit metrics (m³/ton).




Figure 2. Component-Wise Water-Footprint Calculation and Product-Level Output

Validation and Quality Control

Automated validation algorithms inspect incoming data records for missing fields, physical unit mismatches, out-of-bound parameter values, and logical inconsistencies between inputs and outputs. Quality flags (e.g., Validated, Pending Review, and Anomalous) are assigned to ensure high data integrity prior to reporting.

Proposed System Architecture

The software architecture connects data acquisition layers, relational/document database systems, automated calculation engines, and interactive reporting dashboards into a unified traceable system.

Results and evaluation framework Proposed System Output

The primary output of the system is an audit-ready, standardized digital record containing disaggregated green, blue,

and grey water components, total footprint per ton, data source attributions, and quality control flags.

Traceability and Provenance

Traceability is maintained by binding calculated results with raw source inputs, mathematical formulas, model version identifiers, and immutable system timestamps stored within the digital record.

Sequential Workflow Analysis

The sequential data workflow from field data collection by farmers/researchers to dashboard reporting is illustrated in Figure 3.




Figure 3. Sequential Workflow from Agricultural Data Collection to Reporting

Evaluation Framework & Performance Benchmarks

To evaluate system effectiveness, a quantitative evaluation matrix assessing data completeness, validation accuracy, calculation consistency, traceability score, and usability was defined. Figure 4 contrasts the target performance metrics of the proposed digital framework against traditional paper/manual record keeping systems.




Figure 4. Evaluation Framework - Target Performance & Validation Benchmarks

DISCUSSION

System Interpretation

Digitizing water footprint accounting resolves the historical challenge of fragmented, non-traceable agricultural data. By disaggregating green, blue, and grey water components, the framework delivers actionable insights for sustainable resource allocation and product eco-labeling.

Role of Digital Technologies & Smart Agriculture

Digital databases ensure immutable data preservation, while interactive dashboards enable rapid cross-commodity

comparison. Automated integration with IoT soil moisture sensors and satellite remote sensing further reduces human data collection overhead.

Limitations

System accuracy remains dependent on input data resolution and localized weather station availability. Variances in regional water quality standards can also impact grey water footprint calculations.

Future Scope

Future work will focus on deploying cloud-native micro services, building real-time API connectors for IoT smart irrigation networks, and integrating machine learning algorithms for predictive water footprint forecasting under climate change scenarios.

CONCLUSION

This research has presented a robust digital technology framework for calculating, validating, organizing, and reporting water footprints for agricultural products. By combining disaggregated green-blue-grey water calculation engines with structured digital schemas, automated validation, and interactive dashboards, the system achieves unprecedented data traceability and transparency. The proposed architecture serves as a scalable foundation for sustainable agricultural water management, eco-labeling certification, and future smart agriculture integration.

REFERENCES

- [1] M. M. Mekonnen and A. Y. Hoekstra, "The green, blue and grey water footprint of crops and derived crop products," *Hydrology and Earth System Sciences*, vol. 15, no. 5, pp. 1577–1600, 2011.
- [2] A. Y. Hoekstra, A. K. Chapagain, M. M. Mekonnen, and R. E. Aldaya, *The Water Footprint Assessment Manual: Setting the Global Standard*, Earthscan, London/Washington, DC, 2011.
- [3] International Organization for Standardization, "ISO 14046:2014: Environmental management — Water footprint — Principles, requirements and guidelines," ISO, Geneva, 2014.
- [4] M. M. Mekonnen and A. Y. Hoekstra, "A global and high-resolution assessment of the green, blue and grey water footprint of wheat," *Hydrology and Earth System Sciences*, vol. 14, no. 7, pp. 1259–1276, 2010.
- [5] R. G. Allen, L. S. Pereira, D. Raes, and M. Smith, *Crop Evapotranspiration: Guidelines for Computing Crop Water Requirements*, FAO Irrigation and Drainage Paper 56, FAO, Rome, 1998.
- [6] C. Parra-López, S. Sayadi, and J. A. Carmona, "Digital technologies for water use and management in agriculture: Recent applications and future outlook," *Agricultural Water Management*, vol. 309, p. 109347, 2025.
- [7] A. Kumar, S. Sharma, and R. Patel, "Artificial intelligence and IoT for water saving in agriculture: A systematic review," *Smart Agricultural Technology*, vol. 11, p. 101008, 2025.
- [8] V. Gupta, P. Singh, and M. Verma, "Internet of Things-enabled smart irrigation systems for precision water management: A systematic review," *Agricultural Water Management*, vol. 333, p. 110615, 2026.
- [9] Food and Agriculture Organization of the United Nations, *The State of the World's Land and Water Resources for Food and Agriculture: Managing Systems at Risk*, FAO, Rome, 2011.
- [10] International Organization for Standardization, "ISO/TR 14073:2017: Environmental management — Water footprint— Illustrative examples on how to apply ISO 14046," ISO, Geneva, 2017.